

Asthma Action Plan



General Information:

Name _____
 Emergency contact _____ Phone numbers _____
 Physician/Health Care Provider _____ Phone numbers _____
 Physician Signature _____ Date _____

| Severity Classification | Triggers | Exercise |
|--|---|--|
| <input type="radio"/> Mild Intermittent <input type="radio"/> Moderate Persistent <input type="radio"/> Mild Persistent <input type="radio"/> Severe Persistent | <input type="radio"/> Colds <input type="radio"/> Smoke <input type="radio"/> Weather <input type="radio"/> Exercise <input type="radio"/> Dust <input type="radio"/> Air pollution <input type="radio"/> Animals <input type="radio"/> Food <input type="radio"/> Other _____ | 1. Pre-medication (how much and when) _____ 2. Exercise modifications _____ |

Green Zone: Doing Well

Peak Flow Meter Personal Best = _____

Symptoms

- Breathing is good
- No cough or wheeze
- Can work and play
- Sleeps all night

Control Medications

| Medicine | How Much to Take | When To Take It |
|----------|------------------|-----------------|
| _____ | _____ | _____ |
| _____ | _____ | _____ |
| _____ | _____ | _____ |

Peak Flow Meter

More than 80% of personal best or _____

Yellow Zone: Getting Worse

Contact Physician if using quick relief more than 2 times per week.

Symptoms

- Some problems breathing
- Cough, wheeze or chest tight
- Problems working or playing
- Wake at night

Continue control medicines and add:

| Medicine | How Much to Take | When To Take It |
|----------|------------------|-----------------|
| _____ | _____ | _____ |
| _____ | _____ | _____ |
| _____ | _____ | _____ |

Peak Flow Meter

Between 50 to 80% of personal best or _____ to _____

IF your symptoms (and peak flow, if used) return to Green Zone after one hour of the quick relief treatment, THEN

- Take quick-relief medication every 4 hours for 1 to 2 days
- Change your long-term control medicines by _____
- Contact your physician for follow-up care

IF your symptoms (and peak flow, if used) DO NOT return to the GREEN ZONE after 1 hour of the quick relief treatment, THEN

- Take quick-relief treatment again
- Change your long-term control medicines by _____
- Call your physician/Health Care Provider within _____ hours of modifying your medication routine

Red Zone: Medical Alert

Ambulance/Emergency Phone Number: _____

Symptoms

- Lots of problems breathing
- Cannot work or play
- Getting worse instead of better
- Medicine is not helping

Continue control medicines and add:

| Medicine | How Much to Take | When To Take It |
|----------|------------------|-----------------|
| _____ | _____ | _____ |
| _____ | _____ | _____ |
| _____ | _____ | _____ |

Peak Flow Meter

Between 0 to 50% of personal best or _____ to _____

Go to the hospital or call for an ambulance if

- Still in the red zone after 15 minutes
- If you have not been able to reach your physician/health care provider for help
- _____

Call an ambulance immediately if the following danger signs are present

- Trouble walking/talking due to shortness of breath
- Lips or fingernails are blue